

Question 20

- A 23 year of woman presents with low-grade fever, malaise, nocturnal diaphoresis and weight loss over the past 8 weeks. On exam she has a weak right radial pulse and a left infra-scapular bruit. In the last week she has had bilateral transient visual blurring. Her BP=184/50 mm Hg. Her arch angiogram is shown below. Which of the following options is most applicable to her clinical condition:

- A. After ESR, VDRL, ANA, coagulation and serum chemistries she should undergo prompt endovascular intervention
- B. Immediate control of her BP can be deleterious
- C. Treatment with corticosteroids and cyclophosphamide is needed
- D. Angioplasty offers durable success
- E. Regardless of the treatment, she faces a low relapse rate



Question 20

- A 23 year of woman presents with low-grade fever, malaise, nocturnal diaphoresis and weight loss over the past 8 weeks. On exam she has a weak right radial pulse and a left infra-scapular bruit. In the last week she has had bilateral transient visual blurring. Her BP=184/50 mm Hg. Her arch angiogram is shown below. Which of the following options is most applicable to her clinical condition:

- A. After ESR, VDRL, ANA, coagulation and serum chemistries she should undergo prompt endovascular intervention
- B. Immediate control of her BP can be deleterious
- C. Treatment with corticosteroids and cyclophosphamide is needed
- D. Angioplasty offers durable success
- E. Regardless of the treatment, she faces a low relapse rate

