

# Question 16

- In counseling a patients with RAS, which of the following statements would be appropriate to share with a patient with refractory hypertension on 3 medications?
  - A. Clinical trials of RAS stenting compared to medical therapy alone have not shown any significant reduction in BP nor the preservation of renal function long-term
  - B. Angioplasty is often followed by stenting of FMD typically involving the proximal renal artery
  - C. Ostial RAS is better treated with stenting than angioplasty alone
  - D. A 10% residual or a <math><10\text{ mmHg}</math> gradient suggests successful PTA
  - E. ACEI are contraindicated in patients with unilateral RAS due to the risk of azotemia from a sudden increase in transglomerular pressure gradient



# Question 16

- In counseling a patients with RAS, which of the following statements would be appropriate to share with a patient with refractory hypertension on 3 medications?
  - A. Clinical trials of RAS stenting compared to medical therapy alone have not shown any significant reduction in BP nor the preservation of renal function long-term
  - B. Angioplasty is often followed by stenting of FMD typically involving the proximal renal artery
  - C. Ostial RAS is better treated with stenting than angioplasty alone
  - D. A 10% residual or a <math><10\text{ mmHg}</math> gradient suggests successful PTA
  - E. ACEI are contraindicated in patients with unilateral RAS due to the risk of azotemia from a sudden increase in transglomerular pressure gradient

